

10t 10
11/8/97

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ditrac® Rodenticide

ACTIVE CONSTITUENT: 0.05g/kg BRODIFACOUM

ONE FEED - FAST ACTING
EASY TO USE

2.5 kg NET

49868/0997

N.R.A.
APPROVED LABEL

Date 17/9/97

Max Davison

L 07 10
11/8/97

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ditrac® Rodenticide

ACTIVE CONSTITUENT: 0.05g/kg BRODIFACOUM

ONE FEED - FAST ACTING
EASY TO USE

5 kg NET

**Mfg. By: Bell Laboratories, Inc.
Madison, WI 53704 USA**

0 0 1 1 0
11/8/97

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ditrac[®] Rodenticide

ACTIVE CONSTITUENT: 0.05g/kg BRODIFACOUM

ONE FEED - FAST ACTING
EASY TO USE

10 kg NET

Mfg. By: Bell Laboratories, Inc.
Madison, WI 53704 USA

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ditrac® Rodenticide

ACTIVE CONSTITUENT: 0.05g/kg BRODIFACOUM

ONE FEED - FAST ACTING
EASY TO USE

PRIMARY PACK

CONTAINS 100 x 50g BAIT STATIONS WHICH IT IS ILLEGAL TO SELL
SEPARATELY.
DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS

5kg NET

**Mfg. By: Bell Laboratories, Inc.
Madison, WI 53704 USA**

Australian Agent:

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ditrac® Rodenticide

ACTIVE CONSTITUENT: 0.05g/kg BRODIFACOUM

ONE FEED - FAST ACTING
EASY TO USE

PRIMARY PACK

CONTAINS 100 x 50g BAIT STATIONS WHICH IT IS ILLEGAL TO SELL
SEPARATELY.
DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS

5kg NET

Mfg. By: Bell Laboratories, Inc.
Madison, WI 53704 USA

Australian Agent:

DIRECTIONS FOR USE

DO NOT exceed 3m between bait stations for mice or 9m for rats

DO NOT place bait stations in open Mice require several bait stations, 50g per station (less than 1g will kill a mouse). Rats require large quantities of bait in one or two locations, 50 - 100g per station (4-6g will kill a rat). For effective control continue until feeding ceases.

Baiting Strategy: Eliminate as far as practicable, all other food sources. Bait infested area. Place baits stations under cover. Inspect bait stations daily. Replace eaten baits. If quickly eaten increase number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

GENERAL INSTRUCTIONS

Pellets are not subject to wind dispersal. Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Record the number and locations of bait stations for purpose of replenishment and later removal. Most deaths occur 4 to 7 days after consumption of bait.

PRECAUTION: KEEP OUT OF REACH OF CHILDREN

Keep away from food and water and surfaces which come in contact with food and water.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. DO NOT place baits in locations which are accessible to domestic animals, livestock or birds. The product is restricted to use in and around buildings, (within 2 metres) or enclosed spaces, e.g. drains.

STORAGE AND DISPOSAL: DO NOT contaminate dams, streams, rivers waterways and drains with the bait or its used container. Store in a tightly closed, original container in a dry well ventilated area, as cool as possible out of direct sunlight. Shake empty container onto baiting site. Break, crush or puncture and bury empty container in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.

SAFETY DIRECTIONS:

Poisonous if swallowed. Do not inhale dust. When using and after use: Do not touch the bait, use scoop or measure. If on skin and after each baiting wash thoroughly with soap and water. Vitamin K1 (Phytomenadione) is antidotal.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Vitamin K1 (Phytomenadione) is antidotal.

Additional information is listed in the Material Safety Data Sheet Ref: BEL3

NRA Approval No: 49868/0997

Batch No:

Date of Manufacture:

IN AN EMERGENCY DIAL 131 126 POISONS INFORMATION CENTRE

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ditrac® Rodenticide

ACTIVE CONSTITUENT: 0.05g/kg BRODIFACOUM

ONE FEED - FAST ACTING
EASY TO USE

Before using this product, read safety directions on outer container. This sachet must be returned to outer pack after use or disposed of as per disposal instructions. Sale of this sachet except in outer container is illegal.

50g NET

**Mfg. By: Bell Laboratories, Inc.
Madison, WI 53704 USA**

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ditrac® Rodenticide

ACTIVE CONSTITUENT: 0.05g/kg BRODIFACOUM

ONE-FEED - FAST ACTING
EASY TO USE

Before using this product, read safety directions on outer container. This sachet must be returned to outer pack after use or disposed of as per disposal instructions. Sale of this sachet except in outer container is illegal.

25g NET

**Mfg. By: Bell Laboratories, Inc.
Madison, WI 53704 USA**

9 of 10
11/8/9

DIRECTIONS FOR USE

The Ditrac® 25g pack is a ready to use 'Bait station'. The feeding behaviour of mice is different to that of rats. Rats require a larger quantity of bait. DO NOT exceed 3m between bait stations for mice and 9m for rats. Place bait stations in areas of known rodent activity. DO NOT place bait stations out in the open.

Baiting Strategy: Eliminate as far as practicable, all other food sources. Bait infested area. Place bait stations under cover. Inspect bait stations daily. Replace eaten baits. If quickly eaten increase number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

GENERAL INSTRUCTIONS

Pellets are not subject to wind dispersal. Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Record the number and locations of bait stations for purpose of replenishment and later removal. Most deaths occur 4 to 7 days after consumption of bait.

PRECAUTION: KEEP OUT OF REACH OF CHILDREN

Keep away from food and water and surfaces which come in contact with food and water.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. DO NOT place baits in locations which are accessible to domestic animals, livestock or birds. The product is restricted to use in and around buildings, (within 2 metres) or enclosed spaces, e.g. drains.

STORAGE AND DISPOSAL: DO NOT contaminate dams, streams, rivers waterways and drains with the bait or its used container. Store in a tightly closed, original container in a dry well ventilated area, as cool as possible out of direct sunlight. Shake empty container onto baiting site. Break, crush or puncture and bury empty container in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.

SAFETY DIRECTIONS:

Poisonous if swallowed. Do not inhale dust. When using and after use: Do not touch the bait, use scoop or measure. If on skin and after each baiting wash thoroughly with soap and water. Vitamin K1 (Phytomenadione) is antidotal.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Vitamin K1 (Phytomenadione) is antidotal.

Additional information is listed in the Material Safety Data Sheet Ref: BEL3

NRA Approval No: 49868/0997

Batch No:

Date of Manufacture:

IN AN EMERGENCY DIAL 131 126 POISONS INFORMATION CENTRE