

PEEL HERE

POISON

KEEP OUT OF REACH OF CHILDREN.
READ SAFETY DIRECTIONS BEFORE
OPENING OR USING

X-verminator™

Single feed, lethal dose

Rodent Blocks

ACTIVE CONSTITUENT: 0.05g/kg BRODIFACOUm

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL.

READY TO USE - FAST ACTING BLOCK BAIT FOR USE IN
AND AROUND HOMES, AGRICULTURAL, COMMERCIAL
AND INDUSTRIAL BUILDINGS

READ THOROUGHLY BEFORE OPENING OR USING.

STORAGE AND DISPOSAL

Store in the tightly closed, original container in a dry well ventilated area, as cool as possible out of direct sunlight. Shake empty container onto baiting site. Containers which have held bait should not be used for any other purpose. Dispose of empty containers and unused or untaken bait by burying in an approved landfill. If no landfill is available, bury containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Search for dead rats or mice – burn or bury them. DO NOT place in refuse bins or rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.

SAFETY DIRECTIONS

Poisonous if swallowed. DO NOT inhale dust. DO NOT touch baits. Use scoop or measure. If on skin, and after each baiting, wash thoroughly with soap and water. Vitamin K1 (Phytomenadione) is antidotal.

FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia: 131126 or New Zealand: 0800 764 766. Vitamin K1 (Phytomenadione) is antidotal.

NOTE TO PHYSICIANS AND VETERINARIANS

Brodifacoum is a long-lasting anticoagulant chemical which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and hemorrhage may result. Vitamin K1 (Phytomenadione) should be administered urgently by medical, appropriate para-medical personnel or veterinarians. If administered intravenously the injection must be given slowly.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for X-Verminator Single Feed Lethal Dose Rodent Blocks is available from www.x-verminator.co.nz/msd.

CONDITIONS OF SALE

As no control can be exercised over the methods or conditions under which this product is used, no responsibility or claim, other than those required by statute, will be accepted for any damage or injury whatsoever arising from the storage, handling, application, use or disposal of this product

DIRECTIONS FOR USE IN ALL STATES

Restraints
DO NOT exceed 3 m between bait stations for mice, or 9 m for rats.
DO NOT place bait station in open.
DO NOT apply baits to crops.

Baiting Strategy
Eliminate as far as practicable all alternative food sources. Bait infested area. Place bait stations under cover. Inspect bait stations daily. Replace eaten baits. If quickly eaten, increase the number of bait stations. Continue observation and replenishment until no more bait is taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

Situation	Pest	Rate	Comments	Critical
Agricultural Commercial Domestic Industrial Buildings	Rats (<i>Rattus norvegicus</i>) Mice (<i>Mus musculus</i>)	1 to 3 blocks per bait station depending on level of infestation	Set out baits in places frequented by mice and rats but not more than 2 m away from buildings etc. For rats: Place 1-3 blocks at intervals of 5-9 m in infested areas. For mice: Place single blocks at 2-3 m intervals in infested areas.	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

GENERAL INSTRUCTIONS

Locate bait stations along walls or rafters and in dark, sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Wax blocks are particularly suitable for use in damp situations where risk of contamination should be minimized. Record the number and locations of baits for purpose of replenishment and later removal. Most deaths occur 4 to 7 days after consumption of bait.

PRECAUTION: KEEP OUT OF REACH OF CHILDREN

Keep away from food and water and surfaces which come in contact with food and water.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the bait or used containers. The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. DO NOT place baits in locations which are accessible to domestic animals, livestock or birds. Symptoms may be undetectable in dogs for at least a week after taking the bait. If you suspect a dog has consumed the bait consult your vet immediately for treatment. The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces, eg. drains.

Registered to Animal Control Products

408 Heads Road, Wanganui, New Zealand. 0064 6344 5302

APVMA Approval Number: 63624/1009

X-verminator™

Single feed, lethal dose